

# The Quantum Spin Hall Effect

H. Buhmann

*Physikalisches Institut, Universität Würzburg, Am Hubland, 97074 Würzburg, Germany*

The increasing understanding of topological phases in condensed matter physics, which was initiated by the quantum Hall effect, has inspired the search for other topological states, especially, in the absence of magnetic fields. As one example the quantum spin Hall (QSH) effect has been proposed for systems with time reversal symmetry and spin-orbit interactions.<sup>1,2)</sup> In a two-dimensional system this new state is characterized by an insulating bulk and two counter-propagating helical edge states. These Kramers pairs fulfill time reversal symmetry and account for a quantized conductance and spin currents which propagate without dissipation. It turned out that HgTe-based quantum well (QW) structures are most suitable candidates for a successful experimental realization.<sup>3)</sup>

In this presentation, experimental requirements are discussed which lead to the observation of quantized spin polarized edge channel transport, one of the main signatures of the QSH effect.<sup>4,5)</sup> Experiments will be presented which demonstrate the stability of the quantized conductance and its non-local character.<sup>6)</sup> Furthermore, it is possible to show evidence for the spin polarization of the QSH edge channels in an all-electrical measurement<sup>7)</sup> which demonstrates the potential of the QSH effect for spin injection and detection in spintronics applications.

- 1) C.L. Kane and E.J. Mele, Phys. Rev. Lett **95**, 226801 (2005).
- 2) B.A. Bernevig and S.C. Zhang, Phys. Rev. Lett. **96**, 106802 (2006).
- 3) B.A. Bernevig, T.L. Hughes, and S.C. Zhang, Science 314, 1757 (2006).
- 4) M. König *et al.*, Science **318**, 766 (2007).
- 5) M. König *et al.* Journ. Phys. Soc. Japn. **77**, 031007 (2008).
- 6) A. Roth *et al.*, Science **325**, 294 (2009).
- 7) C. Brüne *et al.*, accepted for publication in Nature Physics.